

Chiral symmetry breaking patterns in the $U_L(n) \times U_R(n)$ meson model

G. Fejős*

Theoretical Research Division, Nishina Center, RIKEN, Wako 351-0198, Japan

Chiral symmetry breaking patterns are investigated in the $U_L(n) \times U_R(n)$ meson model. It is shown that new classes of minima of the effective potential belonging to the center of the Lie-algebra exist for arbitrary flavor number n . The true ground state of the system is searched non-perturbatively and although multiple local minima of the effective potential may exist, it is argued that strictly a $U_L(n) \times U_R(n) \longrightarrow U_V(n)$ spontaneous symmetry breaking is possible. The reason behind this is the existence of a discrete subset of axial-symmetries, which connects various $U_V(n)$ symmetric vacua of the theory. The results are in agreement with the Vafa-Witten theorem of QCD, illustrating that it remains valid, even without gauge fields, for this certain effective model of the strong interaction.

PACS numbers: 11.30.Qc, 11.30.Rd

Keywords: Chiral symmetry breaking

I. INTRODUCTION

Quantum chromodynamics (QCD) displays an approximate global $U_L(3) \times U_R(3)$ chiral symmetry. The observed mesonic spectra shows that, this symmetry is broken spontaneously in the ground state of QCD matter. Around the temperature of the scale of QCD (Λ_{QCD}), strongly interacting matter undergoes a chiral symmetry restoration, accompanied by the quark deconfinement transition [23]. Due to difficulties of solving QCD at low energies, effective models built upon the approximate chiral symmetry can give reasonable results regarding the properties of the chiral transition and therefore the phase diagram of the strongly interacting matter. The n -flavored matrix model displaying a $U_L(n) \times U_R(n)$ symmetry [1] give reasonable results for the mesonic spectra and phase diagram [11–13], and can be extended with quark degrees of freedom (quark-meson model) [14, 15] and/or the Polyakov loop [16–21]. Although these models cannot be solved exactly either, they provide a simpler theoretical framework than QCD to describe the behavior of the strongly interacting matter at finite temperature and density. The $U_A(1)$ anomaly of QCD can be easily included into the model through 't Hooft's determinant term [3], however it is argued that close to the critical point the instanton density causing the anomaly is suppressed exponentially [9]. Furthermore, recently new signs of the restoration of the $U_A(1)$ factor around the critical temperature appeared in the literature for $n = 2$ [29], therefore close to the phase transition the anomaly is not expected to play an important role and the pure $U_L(n) \times U_R(n)$ symmetric model might have phenomenological relevance.

It has been known for a long time that there is no infrared stable fixed point of the renormalization group flow of the $U_L(n) \times U_R(n)$ model in $4 - \epsilon$ ($\epsilon > 0$) dimensions [2]: ϵ expansion shows that if $n \geq 2$, even the existing non-trivial $O(2n^2)$ like fixed point becomes unstable in the direction of the quartic coupling. The non-existence of an infrared stable fixed point indirectly shows a transition, which (if exists) is fluctuation induced, and of first order. This has been confirmed numerically for the $n = 2$ case with the use of the Functional Renormalization Group (FRG) approach [7, 8, 22], and expected to remain true at $n > 2$.

Analysis of the fixed point structure does not tell anything about the symmetry breaking pattern, however, studies of the $U_L(n) \times U_R(n)$ model in the past assumed a priori that the symmetry is broken to $U_V(n)$, or in other words, the vacuum expectation value of the dynamical matrix field is proportional to the unit matrix. Presumably, the reason behind this is the Vafa-Witten theorem [4], which states that in vector-like gauge theories with vanishing θ -angles (e.g. QCD) and non-zero but equal fermion masses, global vector-symmetries can not be broken spontaneously. Since the $U_L(n) \times U_R(n)$ symmetric theory is an effective model of QCD, it is plausible to accept that the original statement remains true, however, it should not be considered correct without further exploration. This is especially true for vanishing quark masses (i.e. no explicit breaking of chiral symmetry), for which the theorem only states that if a vector-symmetry breaking state existed, its energy could not be lower than the symmetry conserving one's. The situation is getting more puzzling, if we accept the results of [24], where (in the large- n limit) a new local minimum of the energy functional realizing a symmetry breaking pattern $U_L(n) \times U_R(n) \longrightarrow U_V(n-1)$ was found, from which the system can undergo a first order transition with appropriately chosen model parameters [25]. Furthermore, in [26, 27] it was conjectured that Yukawa couplings of the lepton masses of the Standard Model might be described by the spontaneous breaking of a hypothetic $SU(3) \times SU(3)$ symmetry, realizing yet another ground state which breaks the symmetry to $SU(2) \times SU(2) \times U(1)$. This solution has already been reported in [5] for a generic number of flavors. The existence of these different vacua in the $U_L(n) \times U_R(n)$ model might be in apparent contradiction with the Vafa-Witten theorem, therefore it is an open question whether these minima are not physically applicable for some reason, or the applied approximations are inconsistent.

*Electronic address: fejos@riken.jp

The goal of this paper is to answer the aforementioned questions and to map the vacuum structure in an extended space of the condensate space, searching for new local minima of the effective potential for arbitrary flavor numbers. The paper is organized in a pedagogical way and goes from the most simple cases to a general analysis. In Sec. II, we introduce the model and the symmetry breaking we would like to study in the first place. We list the group invariants and discuss the possibility of breaking of the diagonal $U_V(n)$ symmetry. In Sec. III, we solve the field equations at classical level, and show the energy properties with the corresponding spectra of the found solutions. In Sec. IV we include quantum fluctuations using Functional Renormalization Group (FRG) formalism. We start with the $n = 2$ case in the so-called Local Potential Approximation (LPA), then generalize the arguments for the exact FRG formulation for $n > 2$. It is followed by an analysis of having a most general condensate, and a discussion of obtaining all the $U_V(n)$ symmetric ground states. Conclusions can be found in Sec. V.

II. THE MODEL AND SPONTANEOUS SYMMETRY BREAKING

The dynamical variable of the $U_L(n) \times U_R(n)$ scalar theory is a matrix field M , as an element of the $U(n)$ Lie-algebra. It can be parametrized as:

$$M = (s^a + i\pi^a)T^a, \quad (a = 0, \dots, n^2 - 1), \quad (1)$$

where s^a and π^a are the scalar and pseudoscalar fields, respectively, with T^a being the group generators satisfying $\text{Tr}(T^a T^b) = \delta^{ab}/2$. The Lagrangian of the model is

$$\mathcal{L} = \partial_\mu M \partial^\mu M^\dagger - m^2 \text{Tr}(MM^\dagger) - \frac{g_1}{n^2} [\text{Tr}(MM^\dagger)]^2 - \frac{g_2}{n} \text{Tr}(MM^\dagger MM^\dagger). \quad (2)$$

In order to have a classical potential bounded from below, we require $g_1 + g_2 > 0$ and $g_2 > 0$ [2]. It is clear that, (2) is invariant under the symmetry transformation

$$M \longrightarrow U_R M U_L^\dagger, \quad (3)$$

U_L and U_R are independent $U(n)$ matrices realizing the chiral symmetry with parameters θ_R^a and θ_L^a . These transformations can be reformulated in terms of vector- and axialvector-symmetries as:

$$M \longrightarrow V M V^\dagger, \quad M \longrightarrow A^\dagger M A^\dagger, \quad (4)$$

where V and A are also unitary matrices with parameters $\theta_{V,A}^a = (\theta_R^a \pm \theta_L^a)/2$, respectively. Even though in QCD the axial symmetry is broken anomalously at quantum level, in this paper we do not investigate quantitatively the role of this anomaly, nor any explicit symmetry breaking. Following [24], first we are interested in the possibility of the symmetry breaking pattern

$$\langle M \rangle = v_0 T^0 + v_8 T^8, \quad (5)$$

which means that we set all the expectation values in advance to zero, and investigate only on the $v_0 - v_8$ plane. We will come to the question of general condensates later. In (5) T^0 is the generator of the $U(1)$ subgroup of $U(n)$ and T^8 refers to the longest diagonal generator, regardless of the actual number of flavors (see also Appendix A). This notation is just an adaption from the phenomenologically most important $n = 3$ case. Note that, pattern (5) does not apply for $n = 2$, since in this case there are only four generators. Besides T^0 , only of them is diagonal: T^3 , the third Pauli matrix. Formulae given below can be applied to $n = 2$ by considering T^3 instead of T^8 , i.e. substitute index '3' instead of '8' everywhere.

In terms of the coefficient fields s^a and π^a , (2) turns into

$$\mathcal{L} = \frac{1}{2} \partial_\mu s^a \partial^\mu s^a + \frac{1}{2} \partial_\mu \pi^a \partial^\mu \pi^a - V(s^a, \pi^a), \quad (6)$$

where

$$V = m^2 I_1(s^a, \pi^a) + \frac{g_1}{n^2} I_1^2(s^a, \pi^a) + \frac{g_2}{n} I_2(s^a, \pi^a), \quad (7)$$

with the use of notations

$$I_1(s^a, \pi^a) \equiv \text{Tr}(MM^\dagger) = \frac{1}{2}(s^a s^a + \pi^a \pi^a), \quad (8a)$$

$$I_2(s^a, \pi^a) \equiv \text{Tr}(MM^\dagger MM^\dagger) = \frac{1}{8} d^{abc} d^{aef} (s^b s^c + \pi^b \pi^c)(s^e s^f + \pi^e \pi^f) + \frac{1}{2} f^{abe} f^{acf} s^b s^c \pi^e \pi^f. \quad (8b)$$

Here d^{abc} and f^{abc} are the totally symmetric and antisymmetric tensors of the $U(n)$ group, respectively, see details in

Appendix A. We will need the invariants and their derivatives in a background of v_0, v_8 . The invariants are:

$$I_1|_{v_0, v_8} = \frac{1}{2}(v_0^2 + v_8^2), \quad I_2|_{v_0, v_8} = \frac{1}{4n}(v_0^4 + v_8^4) + \frac{1}{8}(d^{888})^2 v_8^4 + \frac{3}{2n} v_0^2 v_8^2 + \frac{1}{\sqrt{2n}} d^{888} v_8^3 v_0. \quad (9)$$

The needed structure constants are listed in Appendix A, the derivatives of the invariants can be found in Appendix B. The mass matrices are defined as follows:

$$M_{ij}^{2(s)} = \frac{\partial^2 V}{\partial s^i \partial s^j} \Big|_{v_0, v_8}, \quad M_{ij}^{2(\pi)} = \frac{\partial^2 V}{\partial \pi^i \partial \pi^j} \Big|_{v_0, v_8}. \quad (10)$$

At classical level these can be written as

$$M_{ij}^{2(s)} = \left(m^2 + \frac{2g_1}{n^2} I_1|_{v_0, v_8} \right) \frac{\partial^2 I_1}{\partial s_i \partial s_j} \Big|_{v_0, v_8} + \frac{g_2}{n} \frac{\partial^2 I_2}{\partial s_i \partial s_j} \Big|_{v_0, v_8} + \frac{2g_1}{n^2} \frac{\partial I_1}{\partial s_i} \Big|_{v_0, v_8} \frac{\partial I_1}{\partial s_j} \Big|_{v_0, v_8}, \quad (11a)$$

$$M_{ij}^{2(\pi)} = \left(m^2 + \frac{2g_1}{n^2} I_1|_{v_0, v_8} \right) \frac{\partial^2 I_1}{\partial \pi_i \partial \pi_j} \Big|_{v_0, v_8} + \frac{g_2}{n} \frac{\partial^2 I_2}{\partial \pi_i \partial \pi_j} \Big|_{v_0, v_8}. \quad (11b)$$

Checking the detailed expressions of (11a) and (11b) found in Appendix B, it can be seen that we have to diagonalize in the 0–8 indices. This leads to the multiplet structure $1 \oplus 1 \oplus n(n-2) \oplus 2(n-1)$ in each sectors. Note that, when $n = 3$ these groups correspond to η, η' , three pions and four kaons in the pseudoscalar sector, and σ, f_0 singlets with three a_0 's and four κ 's in the scalar sector. In general, the spectrum may show the possibility of a $U_L(n) \times U_R(n) \rightarrow U_V(n-1)$ symmetry breaking. We need to investigate if this indeed realizes in the minimum of the effective potential.

III. NEW CLASSICAL SOLUTIONS OF THE FIELD EQUATIONS

In this section we present the field equations for the condensates v_0, v_8 and search for all possible solutions of them. We note that the formulae below can be applied to $n = 2$ as well, we just have to consider index ‘3’ instead of ‘8’ everywhere. Using (B1), the field equations are:

$$0 = \frac{\partial V}{\partial s^0} \Big|_{v_0, v_8} = m^2 v_0 + \frac{g_1}{n^2} (v_0^2 + v_8^2) v_0 + \frac{g_2}{n^2} \left(v_0^3 + 3v_8^2 v_0 - \frac{n-2}{\sqrt{n-1}} v_8^3 \right), \quad (12a)$$

$$0 = \frac{\partial V}{\partial s^8} \Big|_{v_0, v_8} = m^2 v_8 + \frac{g_1}{n^2} (v_0^2 + v_8^2) v_8 + \frac{g_2}{n^2} \left(3v_0^2 v_8 - 3 \frac{n-2}{\sqrt{n-1}} v_8^2 v_0 + \frac{n^2 - 3n + 3}{n-1} v_8^3 \right). \quad (12b)$$

Being cubic, these coupled equations can be solved analytically and we obtain nine solutions. Since the potential reflects $(v_0, v_8) \longleftrightarrow (-v_0, -v_8)$ symmetry, we have only five inequivalent solutions. Not considering the parity pairs, we get (see also Fig. 1):

$$v_0^{[I]} = 0, \quad v_8^{[I]} = 0, \quad (13a)$$

$$v_0^{[II]} = \sqrt{\frac{-m^2 n^2}{g_1 + g_2}}, \quad v_8^{[II]} = 0, \quad (13b)$$

$$v_0^{[III]} = (n-2) \sqrt{\frac{-m^2}{g_1 + g_2}}, \quad v_8^{[III]} = 2 \sqrt{\frac{-m^2 (n-1)}{g_1 + g_2}}, \quad (13c)$$

$$v_0^{[IV]} = (n-1) \sqrt{\frac{-m^2 n}{g_1 (n-1) + g_2 n}}, \quad v_8^{[IV]} = \sqrt{\frac{-m^2 n (n-1)}{g_1 (n-1) + g_2 n}}, \quad (13d)$$

$$v_0^{[V]} = \sqrt{\frac{-m^2}{g_1/n + g_2}}, \quad v_8^{[V]} = -\sqrt{\frac{-m^2 (n-1)}{g_1/n + g_2}}. \quad (13e)$$

We can immediately see that, for $n = 2$, solutions (13) are symmetric under the interchange of the two condensates. This is due to the fact that at $n = 2$, there is a $v_0 \longleftrightarrow v_8$ (or more appropriately $v_0 \longleftrightarrow v_3$) symmetry of the Lagrangian and of the field equations as well: e.g. solution [V] is the reflection of solution [IV] with respect to the $v_8 = 0$ line (this is not true for $n > 2$).

Solution [V] shows $\langle M \rangle \sim \text{diag}(0, 0, \dots, 1)$, which has already been reported in [5] and particularly for $n = 3$ in [26, 27], where it has been conjectured to be a ground state which may describe lepton mass hierarchy in the Standard Model. We would like to mention however that, in the stability region (i.e. $g_1 + g_2 > 0, g_2 > 0$) this solution always describes a saddle point (this can be verified by checking the mass matrices [B3]), and therefore not a local minimum of the potential. In general, solutions [II] and [III] are local minima, [I] is a local maximum, the others are saddle points.




FIG. 1: Solutions of field equations (12) for arbitrary $n \geq 2$. As n increases, the solutions (except [I]) get away from the origin. Solution [I] is a local maximum, [II] and [III] are local minima, [IV] and [V] are saddle points.

We note that, if the couplings satisfied $g_2 < 0$ and $g_1 + ng_2 > 0$, the only local minimum would be solution [V], but since in this case the potential is not bounded from below, we do not investigate it further.

In the large- n limit, solutions [II]-[III]-[IV] (after rescalings $v_0 \rightarrow nv_0$, $v_8 \rightarrow \sqrt{n}v_8$) exactly reproduce the three local extrema found in [24]. In this case solutions of v_0 become degenerate, and a picture of a double-well potential formes in the $v_8 > 0$ direction. The appearing new local minimum (solution [III]) was a candidate in [25] to describe the ground state and a first order transition of the system, see also Fig. 2. Our general analysis shows that this structure is although distorted at finite n , it still exists: a new local minimum (solution [III]) emerges for every each n value.

The energies of the solutions are as follows:

$$\begin{aligned} V|_{v_0^{[I]}, v_8^{[I]}} &= 0, & V|_{v_0^{[II]}, v_8^{[II]}} &= -\frac{m^4 n^2}{4(g_1 + g_2)}, & V|_{v_0^{[III]}, v_8^{[III]}} &= -\frac{m^4 n^2}{4(g_1 + g_2)}, \\ V|_{v_0^{[IV]}, v_8^{[IV]}} &= -\frac{m^4 n^2 (n-1)}{4n(g_1 + g_2) - 4g_1}, & V|_{v_0^{[V]}, v_8^{[V]}} &= -\frac{m^4 n^2}{4(g_1 + ng_2)}, \end{aligned} \quad (14)$$

and the following relations hold between them (using obvious shorthand notations):

$$V_{[II]} = V_{[III]} < V_{[IV]} \leq V_{[V]} < V_{[I]}. \quad (15)$$

Equality between $V_{[IV]}$ and $V_{[V]}$ is only possible for $n = 2$, as it should. We see that having two minima with exactly the same energy is a general property for arbitrary n (it is interesting to mention that, the energy over the number of degrees of freedom is even n -independent in these minima). This could make us suspicious, whether there is a symmetry forcing these minima to have the same amount of energy. Let us check the spectrum in these minima as well. First we calculate the masses which belong to [II]:

$$M_{\eta, [II]}^2 = 0, \quad M_{\eta', [II]}^2 = 0, \quad M_{\pi, [II]}^2 = 0, \quad M_{K, [II]}^2 = -m^2 \frac{2g_2}{g_1 + g_2}, \quad (16a)$$

$$M_{\sigma, [II]}^2 = -m^2, \quad M_{f_0, [II]}^2 = -m^2 \frac{2g_2}{g_1 + g_2}, \quad M_{a_0, [II]}^2 = -m^2 \frac{2g_2}{g_1 + g_2}, \quad M_{\kappa, [II]}^2 = 0, \quad (16b)$$

and then the ones corresponding to [III]:

$$M_{\eta, [III]}^2 = 0, \quad M_{\eta', [III]}^2 = 0, \quad M_{\pi, [III]}^2 = 0, \quad M_{K, [III]}^2 = 0, \quad (17a)$$

$$M_{\sigma, [III]}^2 = -m^2, \quad M_{f_0, [III]}^2 = -m^2 \frac{2g_2}{g_1 + g_2}, \quad M_{a_0, [III]}^2 = -m^2 \frac{2g_2}{g_1 + g_2}, \quad M_{\kappa, [III]}^2 = -m^2 \frac{2g_2}{g_1 + g_2}. \quad (17b)$$

It is interesting to note that only excitation multiplicities depend on n , masses do not. We see that (17) describes the usual $U_L(n) \times U_R(n) \rightarrow U_V(n)$ breaking, but the spectrum belonging to (16) is the same! We just have to interchange $K \longleftrightarrow \kappa$, but apart from parity properties that is just a matter of naming the excitations. Since the energies of these states, and also the spectra are the same, both describe the very same symmetry breaking, there is no physical




FIG. 2: Potential energy at $n = \infty$ as a function of v_8 after rescaling $v_8 \rightarrow \sqrt{n}v_8$. The value of v_0 is set to where local minima were found. The energy is shifted by a constant to take its zero at $v_8 = 0$. Surprisingly, the two minima describes the same symmetry breaking pattern.

distinction between them. Therefore, we can conclude that, at least at classical level we found two physically equivalent states, and it cannot be distinguished, which one is the ground state. At this point, we would like to emphasize that, in the large- n limit it can be very misleading that, extrema [II]-[III]-[IV] lie on the same line of $v_0/n = \sqrt{-m^2/(g_1 + g_2)}$ and while minimum [II] has $v_8 = 0$, [III] shows $v_8 \neq 0$ (Fig. 2). Since the value of v_0 is the same for [II] and [III], one would expect that obviously these correspond to different symmetry breaking patterns, but eventually it turns out not to be true.

IV. QUANTUM LEVEL

We would like to make connections with previous works, therefore we choose to include fluctuations in the framework of the Functional Renormalization Group (FRG) formalism [6]. In FRG, one derives a flow equation for a one parameter (k) quantum effective action Γ_k , which interpolates between the classical action (describing micro-physics) and the quantum effective action including fluctuations. For some general bosonic field(s) Φ , the one parameter Schwinger functional in euclidean space is defined as

$$e^{-W_k[J]} = \int \mathcal{D}\Phi e^{-S[\Phi] - \frac{1}{2} \int \Phi \cdot R_k \cdot \Phi - \int J \cdot \Phi}, \quad (18)$$

where J is a source and R_k is a regulator function suppressing fluctuations with momenta lower than k . The k dependent quantum effective action is defined as:

$$\Gamma_k[\bar{\Phi}] = W_k[J] - \int J \cdot \bar{\Phi} - \frac{1}{2} \int \bar{\Phi} \cdot R_k \cdot \bar{\Phi}, \quad (19)$$

which obeys the so-called flow equation:

$$\frac{\partial \Gamma_k}{\partial k} = \frac{1}{2} \text{Tr} \left[\frac{\partial R_k}{\partial k} (\Gamma_k^{(2)} + R_k)^{-1} \right], \quad (20)$$

where the trace has to be taken in both functional and matrix sense, and $\Gamma_k^{(2)}$ is the second functional derivative of Γ_k . The flow starts from a UV cutoff scale Λ , where by definition (S being the classical action)

$$\Gamma_{k=\Lambda} = S, \quad (21)$$

and goes to zero including all the fluctuations, eventually reaching the quantum effective action $\Gamma_{k=0} \equiv \Gamma$.

A common approximation scheme is the Local Potential Approximation (LPA), where it is assumed that Γ_k has the

following form:

$$\Gamma_k[\Phi] = \int_x (\partial_i \Phi(x) \partial_i \Phi(x) + U_k(x; \Phi)), \quad (22)$$

where U_k is the k -dependent effective potential. This is expected to be a good approximation if the anomalous dimension remains small. With the choice of the commonly used regulator function introduced by Litim [10]: $R_k(q) = (k^2 - q^2)\Theta(k^2 - q^2)$, from (20), in 3+1 dimensions we get

$$\frac{\partial U_k}{\partial k} = \frac{k^4}{6\pi^2} \sum_i \frac{1}{k^2 + M_i^2}, \quad (23)$$

where M_i are the eigenvalues of the mass matrix $\partial^2 U_k / \partial \Phi_i \partial \Phi_j$.

A. Analysis of the $n = 2$ case

We start to analyze the quantum effects in LPA at $n = 2$, which is the simplest case. Since the $U(2)$ group has only two independent invariant tensors (i.e. the previously introduced I_1 and I_2), the k -dependent effective potential is a two-variable function: $U_k = U_k(I_1, I_2)$. Using Appendix A, the mass matrices can be easily built up. In general we have:

$$M_{ij}^{2(s)} = \partial_1 U_k \frac{\partial^2 I_1}{\partial s^i \partial s^j} + \partial_2 U_k \frac{\partial^2 I_2}{\partial s^i \partial s^j} + \partial_{11} U_k \frac{\partial I_1}{\partial s^i} \frac{\partial I_1}{\partial s^j} + \partial_{22} U_k \frac{\partial I_2}{\partial s^i} \frac{\partial I_2}{\partial s^j} + \partial_{12} U_k \left[\frac{\partial I_1}{\partial s^i} \frac{\partial I_2}{\partial s^j} + \frac{\partial I_1}{\partial s^j} \frac{\partial I_2}{\partial s^i} \right], \quad (24a)$$

$$M_{ij}^{2(s)} = \partial_1 U_k \frac{\partial^2 I_1}{\partial \pi^i \partial \pi^j} + \partial_2 U_k \frac{\partial^2 I_2}{\partial \pi^i \partial \pi^j} + \partial_{11} U_k \frac{\partial I_1}{\partial \pi^i} \frac{\partial I_1}{\partial \pi^j} + \partial_{22} U_k \frac{\partial I_2}{\partial \pi^i} \frac{\partial I_2}{\partial \pi^j} + \partial_{12} U_k \left[\frac{\partial I_1}{\partial \pi^i} \frac{\partial I_2}{\partial \pi^j} + \frac{\partial I_1}{\partial \pi^j} \frac{\partial I_2}{\partial \pi^i} \right]. \quad (24b)$$

With the help of (B1) and (B2), we can evaluate these matrices at (v_0, v_3) , and after diagonalization in the $0 - 3$ sectors, we recover the results of [28]: the right hand side of (23) indeed depends only on I_1 and I_2 , as it should. The most important observation is that I_1 and I_2 are invariant under the transformation $v_0 \longleftrightarrow v_3$, therefore so does the right hand side of (23). If we start to flow the effective potential, this leads to the observation that since $U_{k=\Lambda}$ itself has $v_0 \longleftrightarrow v_3$ symmetry, it is preserved throughout the flow to $k = 0$. In other words, on the two-dimensional plane (v_0, v_3) , the line $v_3 = 0$ can be exactly mapped onto the line $v_0 = 0$, which leads us to the fact that, the two local minima of (13) evolve along these lines exactly the same way, being undistinguishable physically even at the very end of the flow (i.e. the quantum effective action). This concludes that even though we observed a potential symmetry breaking scenario $U_L(n) \times U_R(n) \longrightarrow U_V(n-1)$, at least at $n = 2$ this turns out to be $U_L(n) \times U_R(n) \longrightarrow U_V(n)$ even at quantum level. Note that, in [7, 8, 22] it is shown numerically that, the system undergoes a first order transition starting from the minimum of the line $v_3 = 0$, therefore it does so from the minimum of the line $v_0 = 0$.

Since we used LPA, and restricted ourselves to the $n = 2$ case, we have not yet reached our final goal: we need to analyze the situation carefully for $n > 2$ without having any approximation. The previous analysis however can easily be generalized, leading us to an exact result.

B. Analysis of the $n > 2$ case

We start from the exact flow equation (20). The right hand side is $U(n) \times U(n)$ invariant by construction, but this time we have more than two invariants, which will be introduced by fluctuations. Since we are interested in the effective potential at homogeneous field configurations, we do not have to care about invariant terms containing field derivatives. The terms which are to appear and relevant to our interest in the effective action are the following traces:

$$\text{Tr}(MM^\dagger), \quad \text{Tr}(MM^\dagger MM^\dagger), \quad \text{Tr}(MM^\dagger MM^\dagger MM^\dagger), \quad \dots \quad (25)$$

Depending on the value n , we have various numbers of independent invariants evolving throughout the flow. The important observation is that all of these are made out of traces of products of $M^\dagger M$. Let us take a closer look at this entity in a background v_0, v_8 . Since generators T^0 and T^8 are real, $M|_{v_0, v_8} = M^\dagger|_{v_0, v_8}$, therefore

$$MM^\dagger|_{v_0, v_8} = (v_0 T^0 + v_8 T^8)^2 = \left[\sqrt{\frac{1}{2n}} \begin{pmatrix} 1 & 0 & \dots & \dots \\ 0 & 1 & & \\ \cdot & & 1 & \\ \cdot & & & \cdot \\ \cdot & & & & 1 \end{pmatrix} v_0 + \sqrt{\frac{1}{2n(n-1)}} \begin{pmatrix} 1 & 0 & \dots & \dots & \cdot \\ 0 & 1 & & & \\ \cdot & & 1 & & \\ \cdot & & & \cdot & \\ \cdot & & & & -(n-1) \end{pmatrix} v_8 \right]^2 \quad (26)$$

$$= \frac{1}{2n} \text{diag} \left(\left(v_0 + \frac{v_8}{\sqrt{n-1}} \right)^2, \left(v_0 + \frac{v_8}{\sqrt{n-1}} \right)^2, \dots, (v_0 - \sqrt{n-1} v_8)^2 \right). \quad (27)$$

The important observation here is that, there are two choices to obtain $MM^\dagger|_{v_0, v_8} \sim \mathbf{1}$, which obviously describes a symmetry breaking $U_L(n) \times U_R(n) \rightarrow U_V(n)$:

$$v_0 + \frac{v_8}{\sqrt{n-1}} = \pm(v_0 - \sqrt{n-1}v_8). \quad (28)$$

The plus sign leads to $v_8 = 0$, which is the obvious choice, but the non-trivial scenario is obtained choosing the minus sign:

$$v_8 = 2\frac{\sqrt{n-1}}{n-2}v_0. \quad (29)$$

(For $n = 2$ we get $v_0 = 0$, as we should.) Checking (13c) (solution [III]), it lies exactly on this line. The existence of that minimum is therefore a consequence of this symmetry, which will be formulated more precisely in the next section. We see that for every point of the $v_8 = 0$ line can be associated with a point of the line $v_8 = 2\frac{\sqrt{n-1}}{n-2}v_0$ in the sense that $MM^\dagger$ does not change. Because of the right hand side of (20) has to depend only on $MM^\dagger$ (and not M itself), the aforementioned two lines evolve exactly the same during the flow. Furthermore, because of continuity, not just the lines themselves, but also an open neighborhood of them have this property. This leads us to the observation that, along these lines not just the value of the potential, but also the appropriately chosen directional derivatives are the same. That is what we saw before explicitly; at classical level the calculated spectrum was the same, just some directions of the (s^a, π^a) space had to be interchanged. This result shows that the applied heavy scalar approximation of [24, 25] might look reasonable, but obviously hinders the way in which the exact result was obtained: without having the scalar excitations included, the $U_L(n) \times U_R(n)$ invariance is broken in (20).

C. Minima of the effective potential for a general condensate

We can formulate the appearance of the new minimum and the symmetry transformation behind it in a more general way, and argue that, it is a discrete subset of axial symmetries which generates new minima. Let us start from a condensate proportional to the unit matrix: $\langle M \rangle = v_0 T^0$. Since T^0 commutes with everything, it is clear that $U_V(n)$ transformations leave $\langle M \rangle$ invariant, while $U_A(n)$ does not:

$$\langle M \rangle \rightarrow V \langle M \rangle V^\dagger = \langle M \rangle, \quad \langle M \rangle \rightarrow A^\dagger \langle M \rangle A^\dagger = \langle M \rangle (A^\dagger)^2. \quad (30)$$

The effective potential is symmetric under $U_V(n) \times U_A(n)$ rotations, but since vector symmetries leave $\langle M \rangle$ invariant, we need to search for appropriate axial-symmetry transformations to get copies of the $U_V(n)$ symmetric vacuum. With the notation $U = (A^\dagger)^2$, the most general transformation we can apply is $\langle M \rangle \rightarrow \langle M \rangle U$, where U is an arbitrary unitary matrix. This immediately shows that not just the invariant tensors, but $\langle M \rangle \langle M^\dagger \rangle$ itself has to remain unchanged, in agreement with the previous subsection. We would like to search for copies of the $\langle M \rangle$ background, where the condensate lies in the center of the Lie-algebra. This means we search for U matrices for which exist $\{\tilde{v}_a\}$ real numbers fulfilling the following equation:

$$\langle M \rangle U = \sum_{a=\text{diag}} \tilde{v}_a T^a. \quad (31)$$

This can be rewritten as

$$U = \sum_{a=\text{diag}} \alpha_a T^a, \quad (32)$$

where $\alpha_a = \sqrt{2n}\tilde{v}_a/v_0$. This shows that, the intersection of the center of the Lie-algebra and the $U(n)$ group itself in the fundamental representation connects various non-trivial vacua of the effective potential: each of these transformations refer to a copy of the original $U_V(n)$ symmetric minimum. Unitarity implies:

$$\mathbf{1} = UU^\dagger = \sum_{a,b=\text{diag}} \alpha_a \alpha_b T^a T^b. \quad (33)$$

Since $T^a T^b = 2(d^{abc} + if^{abc})T^c$, using that f^{abc} is totally antisymmetric and $T^0 = \frac{1}{\sqrt{2n}}\mathbf{1}$, (33) is equivalent to the conditions

$$\frac{n}{2} = \sum_{a=\text{diag}} \alpha_a \alpha_a, \quad 0 = \sum_{a,b=\text{diag}} \alpha_a \alpha_b d^{abc}, \quad (34a)$$

which come from linear independency of the generators. (34) looks simple, but from a practical point of view, it is better to express unitarity in terms of matrix elements of $UU^\dagger$. Using the definition of diagonal generators given in

Appendix A, from (33) we get:

$$\mathbf{1} = \text{diag} \left(\left[\frac{\alpha_0}{\sqrt{2n}} + \sum_{i=1}^{n-1} \frac{\alpha_i}{\sqrt{2i(i+1)}} \right]^2, \dots, \left[\frac{\alpha_0}{\sqrt{2n}} - \frac{\alpha_j j}{\sqrt{2j(j+1)}} + \sum_{i=1}^{n-j-1} \frac{\alpha_{j+i}}{\sqrt{2(j+i)(j+i+1)}} \right]^2, \dots \right). \quad (35)$$

We just showed the zeroth and the j^{th} element of the diagonal ($j \in \{0, \dots, n-1\}$). Requiring each element to be equal to unity, we get n equations for n variables $\{\alpha_j\}$.

We present a possible method to solve these coupled equations. The $(j-1)^{\text{th}}$ and the j^{th} equation look as

$$\left| \frac{\alpha_0}{\sqrt{2n}} - \frac{\alpha_{j-1}(j-1)}{\sqrt{2j(j-1)}} + \frac{\alpha_j}{\sqrt{2j(j+1)}} + \sum_{i=1}^{n-j-1} \frac{\alpha_{j+i}}{\sqrt{2(j+i)(j+i+1)}} \right| = 1 \quad (36a)$$

$$\left| \frac{\alpha_0}{\sqrt{2n}} - \frac{\alpha_j j}{\sqrt{2j(j+1)}} + \sum_{i=1}^{n-j-1} \frac{\alpha_{j+i}}{\sqrt{2(j+i)(j+i+1)}} \right| = 1 \quad (36b)$$

If $j \neq 0$, we can plug (36b) to (36a), and arrive at

$$\alpha_j \sqrt{\frac{j+1}{2j}} - \alpha_{j-1} \sqrt{\frac{j-1}{2j}} \pm 1 = \pm 1, \quad (j \in \{1, \dots, (n-1)\}), \quad (37)$$

which is a recursion for α_j and where the $\pm$ sign on the left hand side is not arbitrary, it depends on the sign of (36b), i.e. the one in the j^{th} equation. Because of that, it is convenient to start the recursion backwards. We have not yet used the last matrix element of (34). Requiring it to be equal to unity, we get the starting point of the recursion:

$$\frac{\alpha_0}{\sqrt{2n}} - \frac{\alpha_{n-1}}{\sqrt{2n}} \sqrt{n-1} = \pm 1. \quad (38)$$

In practice we start with (38), choose a sign and express α_{n-1} with α_0 . Then we plug the expression of α_{n-1} into (37) (sign on the left hand side has already been determined with the previous choice!) for $j = n-1$, and choose a sign again for the right hand side. The procedure continues until $j = 1$, when α_1 is determined explicitly, and with that, we get the expression of α_0 and therefore all the $\{\alpha_i\}$ immediately.

The solution described in the previous subsection refers to a set of choice, where at each but the first step *plus* sign is chosen: it leads to

$$\tilde{v}_0 = v_0(n-2)/n, \quad \tilde{v}_{n-1} = 2v_0\sqrt{n-1}/n, \quad \tilde{v}_i = 0 \quad (i = 1, \dots, n-2), \quad (39)$$

which is consistent with (29) or (13b) with (13c). (Note that previously index notation “8” was used to $(n-1)$ regardless of the value of n .) Choosing plus signs only leads to the trivial

$$\tilde{v}_0 = v_0, \quad \tilde{v}_i = 0 \quad (i = 1, \dots, n-1) \quad (40)$$

solution.

With the procedure outlined above, since there are 2^n ways of choosing signs, we can find in general all the 2^n solution of (35) very easily, which connects the usual $v_0 \neq 0, v_i = 0$ line with appropriately chosen directions, and which give every local minima of the potential (7) on a diagonal background. By construction, these turn out to be $U_V(n)$ symmetric ground states, and more importantly, evolve identically with the FRG flow. This illustrates that vacua appearing at classical level can not break the $U_V(n)$ symmetry, and because of the discussed invariance of $MM^\dagger$, this property remains true after including quantum fluctuations too. We note however, although it is not expected, but cannot be excluded that, higher order operators may develop even more new local minima not appearing at classical level.

Finally, let us make a comparison between the Vafa-Witten theorem [4] and our results. The theorem claims that, in vector-type gauge theories with zero θ -vacuum angles (e.g. QCD), if the fermion masses are equal and non-zero, none of the vector symmetries can be broken spontaneously. Our result mainly agrees with this, remarkably without the appearance of any gauge fields, in an effective model. Adding the following explicit symmetry breaking term to the Lagrangian ($H = h_a T^a/4$):

$$\mathcal{L}_h = \text{Tr} (H(M + M^\dagger)) \equiv h_a s^a, \quad (41)$$

paper [4] proves that if $H \sim \mathbf{1}$, which refer to equal quark masses, pattern of the symmetry breaking is $U_L(n) \times U_R(n) \longrightarrow U_V(n)$, moreover, it claims that, if $H = 0$ and because of some reason new, maybe even vector-type symmetry breaking minima existed, their energy cannot be lower than the energy of the symmetry conserving state. Our analysis extends this argument and excludes the possibility of having a $U_V(n)$ breaking state. In the chiral limit we do have multiple ground states as [4] suggests it as a possibility, but always the pattern $U_L(n) \times U_R(n) \longrightarrow U_V(n)$ occurs. However,

in general, explicit symmetry breaking may split these states, therefore $H \neq 0$ can create physically inequivalent stable and/or metastable states. For example $H \sim \mathbf{1}$ with $h_0 > 0$ would clearly select solution [II] (13b) as the true ground state. Phenomenology shows [11] that for QCD $h_0 > 0$, $h_8 < 0$, with $h_3 \approx 0$, which makes (13b) even more favorable energetically. Treating the model as an effective theory of strong interaction, this points out the correctness of investigating the $\langle M \rangle = v_0 T^0$ minimum in the first place, but it is important to stress here is that the vacuum structure of the model shows a much richer structure than it has been investigated before.

V. CONCLUSIONS

We investigated the possibility of having a spontaneous symmetry breaking pattern of the $U_L(n) \times U_R(n)$ meson model other than $U_V(n)$. The Vafa-Witten theorem [4] states that for vector-type gauge theories with vanishing θ -vacuum angles and nonzero but equal fermion-masses, none of the vector-like global symmetries can be broken. Since this applies to QCD and the $U_L(n) \times U_R(n)$ model is an effective theory of the strong interaction, it is common to consider that the expectation value of the dynamical matrix field is $\langle M \rangle \sim \mathbf{1}$. However, we have no reason to believe that, the model obeys a theorem valid for QCD, without further investigations.

Following the route of [24, 25], we considered a more general, $\langle M \rangle = v_0 T^0 + v_8 T^8$ type of symmetry breaking, and started our study at classical level. We found that new symmetry breaking minima of the effective potential appears for arbitrary n , also reproducing the large- n result of [24] and a solution given in [5, 26, 27]. Even though the examined pattern is capable of describing the breaking of vector-symmetries, it turned out that, in the observed minima of the effective potential, all of them were conserved. Furthermore, all the states have the very same energy and spectrum, therefore are physically equivalent to the “usual” minima of the effective potential described by the condensate $\langle M \rangle \sim \mathbf{1}$.

We checked if the structure remained the same after including quantum fluctuations, with the use of the Functional Renormalization Group approach. We started with an approximate solution, the Local Potential Approximation for $n = 2$, and found that fluctuations did not change the structure had been described before. Then, generalizing the applied arguments, we showed that the results hold even without any approximations, for arbitrary flavor number n . The reason behind this turned out to be the existence of a discrete subset of the axial-symmetry, which connects several $U_V(n)$ symmetric vacua. This subset can be formulated as the intersection of the center of the $U(n)$ Lie algebra and the group itself. Using this symmetry, we managed to show how to obtain all the possible vacua of the theory considering the most general diagonal condensate. These minima of the effective potential are $U_V(n)$ symmetric by construction, which illustrates the validity of the Vafa-Witten theorem for the $U_L(n) \times U_R(n)$ meson model.

Acknowledgements

The author is grateful to T. Hatsuda, A. Patkós and Zs. Szép for useful suggestions and their careful reading of the manuscript. The author also thanks Y. Hidaka and Y. Tanizaki for illuminating discussions. This work was supported by the Japan Society for the Promotion of Science under ID No. P11795 and by the Foreign Postdoctoral Research program of RIKEN.

Appendix A. PROPERTIES OF THE U(N) GROUP

The $U(n)$ group has n independent diagonal and $n(n-1)$ independent non-diagonal generators leading to n^2 in total. They are traceless and normalized as $\text{Tr}(T^a T^b) = \delta^{ab}/2$. We introduce the generalized Gell-Mann matrices as $\lambda^i = 2T^i$, where T^i are the group generators. The diagonal ones read as:

$$\lambda^{(0)} = \sqrt{\frac{2}{n}} \begin{pmatrix} 1 & & & \\ & 1 & & \\ & & \dots & \\ & & & 1 \end{pmatrix}, \quad \lambda^{(1)} = \begin{pmatrix} 1 & & & \\ & -1 & & \\ & & 0 & \\ & & & \dots \end{pmatrix}, \quad \dots \quad \lambda^{(n-1)} = \sqrt{\frac{2}{n(n-1)}} \begin{pmatrix} 1 & & & \\ & 1 & & \\ & & \dots & \\ & & & -(n-1) \end{pmatrix}. \quad (\text{A1})$$

We note that, except for the last subsection of section IV, throughout the paper conventionally “8” index goes to $\lambda^{(n-1)}$, independently of the actual value of the flavor number n :

$$2T^8 = \lambda^8 \equiv \lambda^{(n-1)}. \quad (\text{A2})$$

The non-diagonal generators form two groups, as generalizations of (x) and (y) type of Pauli matrices in the following sense:

$$(\lambda^{(x,jk)})_{ab} = \delta_{ak}\delta_{bj} + \delta_{aj}\delta_{bk}, \quad (\lambda^{(y,jk)})_{ab} = i\delta_{ak}\delta_{bj} - i\delta_{aj}\delta_{bk}. \quad (\text{A3})$$

Here compact index notations (x, jk) and (y, jk) ($j < k$) were introduced. The structure constants of the algebra are defined through the relation

$$\lambda^a \lambda^b = (d_{abc} + i f_{abc}) \lambda^c, \quad (\text{A4})$$

where d_{abc} is totally symmetric and f_{abc} is totally antisymmetric in their indices. We have:

$$[\lambda^a, \lambda^b] = 2i f_{abc} \lambda^c \quad \implies \quad \text{Tr} [\lambda^a, \lambda^b] \lambda^c = 4i f_{abc}, \quad (\text{A5a})$$

$$\{\lambda^a, \lambda^b\} = 2d_{abc} \lambda^c \quad \implies \quad \text{Tr} [\{\lambda^a, \lambda^b\} \lambda^c] = 4d_{abc}, \quad (\text{A5b})$$

where $[\cdot, \cdot]$ and $\{\cdot, \cdot\}$ refer to commutation and anticommutation, respectively. Alternatively, we can write

$$f_{abc} = \frac{1}{2} \Im \text{Tr} (\lambda^a \lambda^b \lambda^c), \quad d_{abc} = \frac{1}{2} \Re \text{Tr} (\lambda^a \lambda^b \lambda^c). \quad (\text{A6})$$

The structure constants needed for obtaining the mass matrices are:

$$\begin{aligned} d^{0ab} &= \sqrt{\frac{2}{n}} \delta^{ab}, \quad d^{888} = (2-n) \sqrt{\frac{2}{n(n-1)}}, \quad d^{8ij \neq 0,8} = \begin{cases} (2-n) \sqrt{\frac{1}{2n(n-1)}} \delta^{ij}, & \text{if } i, j \in \{(x, jn), (y, jn)\} \\ \sqrt{\frac{2}{n(n-1)}} \delta^{ij}, & \text{else} \end{cases}, \\ d^{a88} &= \sqrt{\frac{2}{n}} \delta^{a0} + d^{888} \delta^{a8}, \quad f^{u8v} = \sqrt{\frac{n}{2(n-1)}} (\delta_{u,(y,jn)} \delta_{v,(x,jn)} - \delta_{u,(x,jn)} \delta_{v,(y,jn)}). \end{aligned} \quad (\text{A7})$$

Appendix B. CLASSICAL MASS MATRICES

In this section we show the exact form of the mass matrices at classical level. First we list the first and second derivatives of the invariants on a background v_0, v_8 needed for the expressions of the masses. The first derivatives:

$$\left. \frac{\partial I_1}{\partial s^i} \right|_{v_0, v_8} = v_0 \delta^{i0} + v_8 \delta^{i8}, \quad \left. \frac{\partial I_2}{\partial s^i} \right|_{v_0, v_8} = \frac{1}{n} v_0^3 \delta^{i0} + \frac{3}{\sqrt{2n}} d^{888} v_8^2 v_0 + \frac{3}{n} v_0^2 v_8 \delta^{i8} + \frac{1}{2} d^{888} d^{i88} v_8^3 + \frac{1}{n} v_8^3 \delta^{8i}, \quad (\text{B1a})$$

$$\left. \frac{\partial I_2}{\partial \pi^i} \right|_{v_0, v_8} = 0, \quad \left. \frac{\partial I_2}{\partial \pi^i} \right|_{v_0, v_8} = 0. \quad (\text{B1b})$$

The non-zero second derivatives:

$$\left. \frac{\partial^2 I_1}{\partial s^i \partial s^j} \right|_{v_0, v_8} = \delta^{ij}, \quad \left. \frac{\partial^2 I_1}{\partial \pi^i \partial \pi^j} \right|_{v_0, v_8} = \delta^{ij}, \quad (\text{B2a})$$

$$\begin{aligned} \left. \frac{\partial^2 I_2}{\partial s^i \partial s^j} \right|_{v_0, v_8} &= \frac{1}{n} (3v_0^2 + v_8^2) \delta^{ij} + 3\sqrt{\frac{2}{n}} d^{8ij} v_0 v_8 + \frac{1}{2} d^{888} d^{8ij} v_8^2 + \frac{2}{n} (\delta^{i8} \delta^{j8} + \delta^{i0} \delta^{j8}) v_8^2 + (d^{8ii})^2 \delta^{ij \neq 0,8} v_8^2, \\ &\quad + d^{888} \sqrt{\frac{2}{n}} (\delta^{i0} \delta^{j8} + \delta^{j0} \delta^{i8}) v_8^2 + (d^{888})^2 \delta^{i8} \delta^{j8} v_8^2, \end{aligned} \quad (\text{B2b})$$

$$\left. \frac{\partial^2 I_2}{\partial \pi^i \partial \pi^j} \right|_{v_0, v_8} = \frac{1}{n} (v_0^2 + v_8^2) \delta^{ij} + \frac{1}{2} d^{888} d^{8ij} v_8^2 + \sqrt{\frac{2}{n}} d^{8ij} v_8 v_0 + \sum_a (f^{a i8})^2 \delta^{ij} v_8^2. \quad (\text{B2c})$$

Relations (B1) are also needed for the derivation of the field equations, (B2) can only be used for determining the mass matrices. Using the expressions of the structure constants (A7), invariants (9) and their derivatives (B1), (B2)

just listed, the nonzero elements of the mass matrices in a background v_0, v_8 are as follows:

$$M_{ij}^{2(s)} = \begin{cases} m^2 + \frac{1}{n^2} (g_1(3v_0^2 + v_8^2) + 3g_2(v_0^2 + v_8^2)), & i = 0, j = 0 \\ \frac{1}{n^2} (2g_1v_0v_8 + g_2(6v_0v_8 - 3\frac{n-2}{\sqrt{n-1}}v_8^2)), & i = 0, j = 8 \text{ or } i = 8, j = 0 \\ m^2 + \frac{1}{n^2} (g_1(v_0^2 + 3v_8^2) + 3g_2(v_0^2 + \frac{n^2-3n+3}{n-1}v_8^2 - 2\frac{n-2}{\sqrt{n-1}}v_0v_8)), & i = 8, j = 8 \\ m^2 + \frac{1}{n^2} (g_1(v_0^2 + v_8^2) + g_2(3v_0^2 + \frac{n^2-3n+3}{n-1}v_8^2 - 3\frac{n-2}{\sqrt{n-1}}v_0v_8)), & i = j \in \{(x, jn), (y, jn)\} \\ m^2 + \frac{1}{n^2} (g_1(v_0^2 + v_8^2) + 3g_2(v_0^2 + v_8^2 + \frac{2}{\sqrt{n-1}}v_0v_8)), & i = j, \text{ else.} \end{cases} \quad (\text{B3a})$$

$$M_{ij}^{2(\pi)} = \begin{cases} m^2 + \frac{1}{n^2} (g_1 + g_2) (v_0^2 + v_8^2), & i = 0, j = 0 \\ \frac{g_2}{n^2} (2v_0v_8 - \frac{n-2}{\sqrt{n-1}}v_8^2), & i = 0, j = 8 \text{ or } i = 8, j = 0 \\ m^2 + \frac{1}{n^2} (g_1(v_0^2 + v_8^2) + g_2(v_0^2 + \frac{n^2-3n+3}{n-1}v_8^2 - 2\frac{n-2}{\sqrt{n-1}}v_0v_8)), & i = 8, j = 8 \\ m^2 + \frac{1}{n^2} (g_1(v_0^2 + v_8^2) + g_2(v_0^2 + \frac{n^2-n+1}{n-1}v_8^2 - \frac{n-2}{\sqrt{n-1}}v_0v_8)), & i = j \in \{(x, jn), (y, jn)\} \\ m^2 + \frac{1}{n^2} (g_1(v_0^2 + v_8^2) + g_2(v_0^2 + \frac{1}{n-1}v_8^2 + \frac{2}{\sqrt{n-1}}v_0v_8)), & i = j, \text{ else.} \end{cases} \quad (\text{B3b})$$

-
- [1] M. Gell-Mann and M. Levy, *Nuovo Cim.* **16**, 705 (1960).
 - [2] R. D. Pisarski and F. Wilczek, *Phys. Rev. D* **29**, 338 (1984).
 - [3] G. 't Hooft, *Phys. Rev. D* **14**, 3432 (1976).
 - [4] C. Vafa and E. Witten, *Nucl. Phys. B* **234**, 173 (1984).
 - [5] A. J. Paterson, *Nucl. Phys. B* **190**, 188 (1981).
 - [6] C. Wetterich, *Phys. Lett. B* **301**, 90 (1993).
 - [7] J. Berges and C. Wetterich, *Nucl. Phys. B* **487**, 675 (1997).
 - [8] J. Berges, N. Tetradis and C. Wetterich, *Phys. Lett. B* **393**, 387 (1997).
 - [9] T. Schafer and E. V. Shuryak, *Rev. Mod. Phys.* **70**, 323 (1998).
 - [10] D. F. Litim, *Phys. Rev. D* **64**, 105007 (2001).
 - [11] J. T. Lenaghan, D. H. Rischke and J. Schaffner-Bielich, *Phys. Rev. D* **62**, 085008 (2000).
 - [12] D. Roder, J. Ruppert and D. H. Rischke, *Phys. Rev. D* **68**, 016003 (2003).
 - [13] T. Herpay, A. Patkós, Zs. Szép and P. Szépfalusy, *Phys. Rev. D* **71**, 125017 (2005).
 - [14] N. Bilic and H. Nikolic, *Eur. Phys. J. C* **6**, 513 (1999).
 - [15] P. Kovács and Zs. Szép, *Phys. Rev. D* **75**, 025015 (2007).
 - [16] T. Kähära and K. Tuominen, *Phys. Rev. D* **78**, 034015 (2008).
 - [17] H. Mao, J. Jin and M. Huang, *J. Phys. G* **37**, 035001 (2010).
 - [18] G. Markó and Zs. Szép, *Phys. Rev. D* **82**, 065021 (2010).
 - [19] B.-J. Schaefer, M. Wagner and J. Wambach, *Phys. Rev. D* **81**, 074013 (2010).
 - [20] T. K. Herbst, J. M. Pawłowski and B.-J. Schaefer, *Phys. Lett. B* **696**, 58 (2011).
 - [21] V. K. Tiwari, *Phys. Rev. D* **86**, 094032 (2012).
 - [22] K. Fukushima, K. Kamikado and B. Klein, *Phys. Rev. D* **83**, 116005 (2011).
 - [23] K. Fukushima and T. Hatsuda, *Rept. Prog. Phys.* **74**, 014001 (2011).
 - [24] G. Fejős and A. Patkós, *Phys. Rev. D* **84**, 036001 (2011).
 - [25] G. Fejős and A. Patkós, *Phys. Rev. D* **85**, 117502 (2012).
 - [26] E. Nardi, *Phys. Rev. D* **84**, 036008 (2011).
 - [27] J. R. Espinosa, C. S. Fong, E. Nardi, *arXiv:1211.6428*.
 - [28] A. Patkós, *Mod. Phys. Lett. A* **27**, 1250212 (2012).
 - [29] S. Aoki, H. Fukaya and Y. Taniguchi, *arXiv:1209.2061*.